

Essential information
for the supply of
VIAGRA CONNECT[®]

50 mg film-coated tablets (sildenafil)



Introduction	5
SECTION 1	
Understanding erectile dysfunction and VIAGRA CONNECT®	7
SECTION 2	
Who can use VIAGRA CONNECT®?	13
SECTION 3	
What is the VIAGRA CONNECT® Pharmacy Consultation Guide?	19
SECTION 4	
Advice for men who are suitable for VIAGRA CONNECT®	23
SECTION 5	
Patient supply scenarios	29
SECTION 6	
Case Study Responses	35
SECTION 7	
Essential information	37
References	41

INT
ROD
UCT
ION

Introduction

This booklet is part of a package of information and training materials that Pfizer Healthcare Ireland have put in place to support the launch of VIAGRA CONNECT as an OTC treatment for Erectile Dysfunction. This document contains essential information you will need to know and consider when assessing men as suitable candidates for use of VIAGRA CONNECT.

This training can form part of your CPD training and once completed can be recorded on your ePortfolio of your IIOP portal.

The sections in this booklet contain the following information:

SECTION 1

Understanding erectile dysfunction and VIAGRA CONNECT

Provides an understanding of VIAGRA CONNECT and how it works.

SECTION 2

Who can use VIAGRA CONNECT?

Details the types of men who can safely use VIAGRA CONNECT taking into account comorbidities and concomitant drug use.

SECTION 3

What is the VIAGRA CONNECT Pharmacy Consultation Guide?

The VIAGRA CONNECT Pharmacy Consultation Guide is provided as an optional tool to help accurately assess patients and to validate their suitability for treatment.

SECTION 4

Providing advice about VIAGRA CONNECT

Information on the advice you can offer patients who are suitable for VIAGRA CONNECT and follow-up advice for those who are found to be not suitable.

SECTION 5

Patient supply scenarios

Test your knowledge with a number of patient supply scenarios to which you can apply your professional judgement and check against the suggested supply outcomes.

SECTION 6

Case study responses

SECTION 7

Essential information

The essential information from the Summary of Product Characteristics.

Understanding erectile dysfunction and VIAGRA CONNECT®

What is Erectile Dysfunction?

Erectile Dysfunction (ED) has been defined as the inability to attain and/or maintain an erection hard enough for satisfactory sexual intercourse.¹ It can occur occasionally or frequently and men are able to self-recognise the condition, which can vary in severity from mild (a soft erection) to severe (no erection at all).

Causes of ED?

Understanding the causes and conditions associated with ED can help you manage the sufferer appropriately. In general, ED is due to 3 main causes:²

- Physical causes
- Psychological causes
- Some types of medication

1. Physical causes

ED usually has an underlying physical cause, and it is useful to bear this in mind when managing patients who have co-morbidities that may predispose them to ED. These could include hypertension, diabetes mellitus, hypercholesterolaemia or CV disease.³

2. Psychological causes:

Psychological causes inhibit normal stimuli to and from the brain and can interfere with the ability to achieve a normal erection. Although there can be a perception that ED is a psychological issue, the reality is, purely psychological causes account for only 1 in 10 cases.⁴

Psychological causes can include:⁵

- Performance anxiety in relation to ED and the fear that it will keep occurring
- Depression – the link with ED appears to be bidirectional. In depressed men, ED may be a symptom of depression, and in men with ED, the emotional stress commonly associated with loss of sexual function may lead to depression
- Antidepressant drugs, such as the tricyclics and selective serotonin reuptake inhibitors, can also cause ED⁶
- Lack of arousal and/or inhibition between partners
- Other sexual dysfunction (e.g. premature ejaculation)
- Major life stress such as; money worries, bereavement, etc
- Low self-esteem – this can be due to prior episodes of ED (thus a feeling of inadequacy) or can be the result of other issues unrelated to sexual performance
- Indifference – this may come about as a result of age and a subsequent loss of interest in sex, or

SECTION 1: Understanding erectile dysfunction and VIAGRA CONNECT

stem from problems in a couple's relationship

3. Medication-induced

Although some drugs can directly cause erection problems (iatrogenic cause), many of the drugs shown in Table 1 are used to treat diseases that are themselves associated with ED.

Recreational drugs, including alcohol, can also cause erection problems.

Table 1: Medications that can cause ED⁶⁻¹²

What is VIAGRA CONNECT and what is it used for?

Therapy area	Drug class or drug	Impact on ED
Cardiovascular	Diuretics	Not determined, but believed to interfere with smooth muscle relaxation
	ACE and ARB inhibitors	Interferes with smooth muscle relaxation
	Aldosterone antagonists	An anti-adrenergic action
	Beta-blockers	Affects sex hormones, impairs vasodilation of the corpora cavernosa
	Clonidine	Depresses adrenergic output
Psychotropic	Anti-depressants, e.g. selective serotonin reuptake inhibitors, tricyclics, monoamine oxidase inhibitors, lithium	Decreases arousal and desire
	Anti-psychotics, e.g. phenothiazines, butyrophenones	Increases prolactin levels
Anti-epileptics	Carbamazepine Phenytoin Barbiturates	Affects sex hormone levels

Table 1 continued

Therapy area	Drug class or drug	Impact on ED
Endocrine drugs	Testosterone antagonists/ oestrogen agonists Anabolic steroids Luteinising hormone-releasing hormone analogues	Affects androgen receptors, reducing sexual desire
Recreational drugs	Alcohol, heroin, cocaine, marijuana, methadone	Causes vasoconstriction and/or impacts on neurotransmitters in the erectile pathway
Other	H2 antagonists, ranitidine and cimetidine	Increases prolactin levels, reducing sexual desire
	Cytotoxics, e.g. cyclophosphamide, methotrexate	The effect of nausea and general malaise associated with such drugs often diminish libido

VIAGRA CONNECT contains sildenafil 50 mg and is a pharmacy-only medicine to treat erectile dysfunction (ED) in men aged 18 years and over. It is a well-established and well-tolerated treatment and provides men with access without prescription to a legal ED medication and professional health advice from the pharmacy.


As well as improving their erectile function, allowing men to fully engage in intercourse and/or masturbation, successful treatment with sildenafil can also have emotional benefits for many men with ED, helping them to regain their self-esteem, self-confidence and relationship satisfaction. It can improve their overall quality of life and thus positively impact the wider social implications of erection problems, such as depression.¹³

How does VIAGRA CONNECT work?

When a man is sexually stimulated, impulses from the brain reach the cavernous nerve which releases nitric oxide (NO) at nerve endings in the penis. The NO diffuses across the endothelial cells into smooth muscle and stimulates the enzyme, guanylate cyclase, to convert GTP into cyclic GMP (cGMP). Cyclic GMP starts a further cascade of biochemical reactions which results in smooth muscle relaxation and the cavernous bodies fill with blood and become rigid. Cyclic GMP is normally broken down by the enzyme PDE5 to GMP, which terminates the pathway and produces detumescence. VIAGRA CONNECT inhibits the action of this enzyme thus increasing cGMP levels and maintaining these levels for longer. Hence VIAGRA CONNECT helps a man attain and maintain an erection in response to sexual stimulation.¹⁴

In order for VIAGRA CONNECT to be effective, sexual stimulation is required.

Figure 1: VIAGRA CONNECT mode of action



- PDE5 inhibitor augments the normal physiological response to sexual stimulation
- There is no effect in the absence of sexual stimulation

VIAGRA CONNECT Efficacy

VIAGRA CONNECT is an efficacious treatment for ED at the recommended 50 mg OTC dose. The efficacy of sildenafil has been demonstrated extensively in numerous clinical trials involving more than 23,000 men.

In some patients, 50 mg may not be efficacious and a referral to the GP is required as the dose may need to be increased to 100 mg. Patients who have renal or hepatic impairment, concomitant medication such as CYP3A4 inhibitors or alpha blockers or who have issues with tolerability should also be referred to their doctor as a 25 mg dose may be appropriate. The 25 mg and 100 mg doses are only available on prescription.

In fixed dose studies, the proportions of patients reporting that treatment improved their erections were 74% (50 mg).¹⁵ These data show VIAGRA CONNECT is effective in men with ED and with a wide range of concomitant diseases.

Who can use VIAGRA CONNECT[®]?

Only adult men aged 18 years or older who have erectile dysfunction (ED) can use the product.

The active ingredient in VIAGRA CONNECT, sildenafil, is a part of the class of medicines known as Phosphodiesterase 5 Inhibitors (PDE5i) that are recommended as a first line treatment for ED by The European Association of Urology Guidelines.

VIAGRA CONNECT is licenced for packs of 2, 4 and 8 tablets (not all sizes may be marketed).

There are a number of contraindications or special warnings which mean that in certain circumstances the product cannot be used by men with ED, see below.

Who must not use VIAGRA CONNECT?

- **Men aged under 18 years**, as it not indicated for this age group. These men should be directed to their GP for follow up if they believe they have an issue relating to erectile function
- **Women**, as it is only indicated for men aged 18 years and over. For a woman interested in the product for their male partner, it is important to ask them to encourage their partner to visit the pharmacy or their doctor for additional advice
- **Men who do not have ED. It is important that you establish that the man has a problem getting or keeping an erection which is satisfactory for sexual performance.** VIAGRA CONNECT will not enhance men's erections or sexual performance and it will not help problems such as premature ejaculation. Men in the latter group should be directed to their GP for further advice
- **Men allergic to sildenafil** or any other ingredient in the medicine

Men with the following health problems must not use VIAGRA CONNECT:

- **Men with hypotension (<90/50 mm Hg) must not use VIAGRA CONNECT.** This is because the safety of sildenafil has not been studied in these sub-groups of patients, and its use is, therefore, contraindicated
- **Men with previously diagnosed mild, moderate or severe hepatic impairment** (e.g. liver cirrhosis), should be advised to talk with their GP about a suitable starting dose or alternative options for the treatment of ED. Men should be asked the questions included in the Pharmacy Consultation Guide (see Section 3) relating to those diagnosed with liver disease or liver problems. If they are under the care of a doctor for liver problems, they will be aware of this and answer 'yes'. As such, this means that you must not supply the product and refer them to their GP
- **Men with previously diagnosed severe renal impairment.** In most cases, patients with severe renal impairment will have some signs or symptoms of the underlying issues and will be under the care of a GP or renal specialist. A question is included in the Pharmacy Consultation Guide which asks the man whether he is suffering from severe kidney problems. If the answer to this is 'yes', then they should be advised to talk with their doctor about alternative options for the treatment of ED
- **VIAGRA CONNECT must not be used in patients with anatomical deformation of the penis** (such as angulation, cavernosal fibrosis or Peyronie's disease) **or in patients with sickle cell anaemia,**

SECTION 2: Who can be supplied VIAGRA CONNECT?

multiple myeloma or leukaemia. There is an increased risk of priapism with these patients and they should be directed to their GP for further advice

- **Sildenafil is contraindicated in patients who have loss of vision in 1 eye because of non-arteritic anterior ischaemic optic neuropathy (NAION),** regardless of whether this episode was in connection with previous PDE5i exposure or not
- **Men who have an inherited eye disease such as retinitis pigmentosa** (a minority of these patients have genetic disorders of retinal phosphodiesterases). This is because the safety of sildenafil has not been studied in these sub-groups of patients, and its use is therefore contraindicated
- **Men who have any bleeding issues** (e.g. haemophilia) **or suffer from stomach ulcers** must not use VIAGRA CONNECT and should be directed to see their GP
- **Rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption** must not take VIAGRA CONNECT because of the lactose contained within the tablet
- **Men who have cardiovascular issues,** see section below

Men with the following cardiovascular (CV) issues must not use VIAGRA CONNECT:

- **Men who get very breathless or experience chest pains with light or moderate physical activity,** such as walking briskly for 20 minutes or climbing 2 flights of stairs should be referred to their GP¹⁷
- **Men who have been advised against sexual activity because of a CV problem.** This group of men are potentially at higher risk due to the burden of over activity on their heart and as such should be referred back to their GP
- **Men with recent history (in the last 6 months) of stroke or myocardial infarction** are also contraindicated from using VIAGRA CONNECT and will need to follow up with their GP
- **Patients with increased susceptibility to vasodilators include those with left ventricular outflow obstruction (e.g. aortic stenosis), or those with the rare syndrome of multiple system atrophy manifesting as severely impaired autonomic control of blood pressure.** Men with these conditions must not use the product without consulting a doctor
- **Patients previously diagnosed with the following must be advised to consult with their GP before resuming sexual activity:**
 - uncontrolled hypertension
 - hypotension
 - unstable angina
 - moderate to severe valvular disease
 - left ventricular dysfunction
 - hypertrophic obstructive and other cardiomyopathies
 - significant arrhythmias
 - severe cardiac failure

Note: Some low risk cardiovascular patients may be suitable for VIAGRA CONNECT, as long as their doctor has advised they can resume sexual activity: those with **asymptomatic controlled hypertension, mild valvular disease or who have had successful coronary artery bypass grafting, stenting or angioplasty.**

Men taking certain other medicines must not use VIAGRA CONNECT

- **Men taking nitrates (e.g. isosorbide mononitrate/dinitrate and glyceryl trinitrate), nitric oxide donors (e.g. nicorandil) or 'poppers' (e.g. amyl nitrate).** Consistent with its known effects on the NO cGMP pathway, sildenafil has been shown to potentiate the hypotensive effects of nitrates, and it is contraindicated for use with nitric oxide donors, nitrates, amyl nitrite (known as the recreational drug 'poppers'), sodium nitroprusside and nicorandil. These men should be directed to their GP for further advice
- **The co-administration of sildenafil with guanylate cyclase stimulators, such as riociguat, is contraindicated** as it may potentially lead to symptomatic hypotension. Again, these men will need to talk with a doctor for further advice
- **Men taking ritonavir,** a potent CYP3A4 inhibitor used in the treatment of HIV, are contraindicated due to the potential for increased blood levels of sildenafil in these patients. They should consult with their doctor for further guidance
- **Men already being treated with another PDE5i** or a higher or lower dose of sildenafil should not use VIAGRA CONNECT. Men taking 50 mg of sildenafil can use the product providing they meet the criteria for pharmacy supply and do not exceed 50 mg as a daily dose
- **Men taking CYP3A4 inhibitors or alpha-blockers** (see Table 2 for examples). These patients should be advised to speak with their GP about a lower suitable starting dose which is available on prescription

Table 2: Examples of CYP3A4 inhibitors and alpha-blockers

Drug class	Drug name
Antibiotics	Erythromycin, clarithromycin, rifampicin
Antifungals	Itraconazole, ketoconazole
Calcium channel blockers	Diltiazem, verapamil
H2-antagonists	Cimetidine
HIV-protease inhibitors	Amprenavir, fosamprenavir, atazanavir, darunavir, indinavir, lopinavir, ritonavir, saquinavir, tipranavir
Alpha blockers	Phenoxybenzamine, phentolamine, tolazoline, trazodone, alfuzosin, doxazosin, tamsulosin, prazosin, terazosin

Case studies

Case study 1:

How would you respond to the following patient request?

Mr. P is a man in his mid-50s who is new to your pharmacy. He has walked briskly to the pharmacy as it is raining and asks to see you. The first thing he states is: "I get out of breath so quickly these days; I could do with a sit down." He then goes on to tell you he would like to purchase VIAGRA CONNECT.

How would you handle this conversation? Would you sell VIAGRA CONNECT in this situation?

See page 35 for case study responses

Case study 2:

How would you respond to the following patient request?

Mr. G is a 72-year-old man who has asked to talk with you about the 'new VIAGRA'. After questioning, you discover he is a smoker with a history of cardiovascular issues including angina, despite undergoing bypass surgery 3 years ago. He is on a range of cardiac medications including glyceryl trinitrate spray for his chest pains. He also has a number of other issues including type 2 diabetes and benign prostatic hypertrophy treated with tamsulosin. His wife has seen a newspaper advert for the product and reckons it is just what he needs.

How would you handle this conversation? Would you make a sale of VIAGRA CONNECT in this situation?

See page 35 for case study responses

What is the VIAGRA CONNECT[®] Pharmacy Consultation Guide?

The VIAGRA CONNECT Pharmacy Consultation Guide, on pages 20–21 of this document, has been provided as an optional tool to help you when assessing men to accurately validate their suitability for treatment. It also provides information which will enable you to offer valuable health advice for men regardless of whether or not they are suitable for the product.

There is no mandatory requirement to use the Pharmacy Consultation Guide; it is your professional judgement to decide when and how to use as an aid in deciding whether to supply VIAGRA CONNECT to the man. The recommendation, however, is to use the supplied questions as a framework to assess suitability. You may also wish to check the patient's medication record to check their concomitant conditions, such as hepatic disease or renal impairment, and medication intake, such as nitrate use.

The VIAGRA CONNECT Pharmacy Consultation Guide includes an assessment of the man's overall fitness for sex, by determining if he gets out of breath or experiences chest pain when he undertakes physical activity. It is also important to understand if the man has any other medical issues that could mean the product is not suitable for him. These may include cardiovascular problems, such as a recent heart attack or stroke which would be revealed through questioning, so the following should be considered:

- There is a degree of cardiac risk associated with sexual activity, although the risk is small in those with stable CVD
- Checking the man's fitness for sex will help identify his CV risk; those with a low risk can be initiated on VIAGRA CONNECT and reminded to follow up with their GP as soon as possible within 6 months
- It also helps identify men at higher risk of CV problems who would benefit from further investigation by their doctor, ideally as soon as possible

Those patients who do not receive the product should be provided with the reason for non-supply and advised to go to the doctor. A tear off piece is included at the bottom of the Pharmacy Consultation Guide for this purpose and contains a section to provide a written explanation of the reason for non-supply. The patient can then take the slip with them to the doctor to help start the conversation.

VIAGRA CONNECT®

50 mg film-coated tablets (sildenafil)

Pharmacy Consultation Guide

The following has been created as a useful aide-memoire to help determine whether your patient is suitable for Viagra Connect, or whether he should be seen by a doctor for further advice. Use of the Pharmacy Consultation Guide is optional, and you should use your professional judgement to decide when and how to use it. The Essential Information for the Supply of Viagra Connect provides additional background information in relation to the supply of this product.

If the patient has previously been supplied with Viagra Connect, he should be asked if anything has changed with respect to his health status or medicines usage. There is no need to repeat the questions below prior to resupply in that case. Remind the patient to make an appointment with his doctor as soon as he can within the first 6 months of starting to use Viagra Connect, to ensure that their erection problems are not caused by any serious health condition. If any factors have changed, Sections 2-4 should be reviewed again. The repeat supply tear-off slip (below) can also be given to the man.

1. Who is Viagra Connect for?

Viagra Connect is only intended for men 18 years and older who are experiencing erectile dysfunction (ED) (i.e. difficulty in getting and/or maintaining an erection satisfactory for sexual performance). This product must not be supplied to men who do not have ED.

It is important to confirm if the man is already receiving treatment for the condition.

Men currently prescribed 50 mg of sildenafil can be supplied this product if they meet the criteria for pharmacy supply, provided they do not take more than 50 mg daily. If the man is using a different dose of sildenafil or another ED treatment, he should be referred to his doctor.

2. Check patient's cardiovascular (CV) health

If the patient answers **YES** to any of the following: **do not supply the product** and refer to the doctor. If you have any reason to consider, based on physical status, the patient should not be using this product, refer to the doctor.

- Y N Has your doctor advised that you are not fit enough for any physical and/or sexual activity?
- Y N Do you feel very breathless or experience chest pain with light or moderate physical activity, such as walking briskly for 20 minutes or climbing two flights of stairs?
- Y N Have you had a heart attack or stroke within the last 6 months?
- Y N Do you have any other heart problems or are you under a doctor's care for any of the following:
- Low blood pressure or uncontrolled high blood pressure
 - Unstable angina (chest pain), irregular heart beat or palpitations (arrhythmia)
 - A problem with one of the valves in your heart (valvular heart disease)
 - A problem where the heart muscle becomes inflamed and does not work as well as it should (cardiomyopathy)
 - Heart problems causing blood flow issues (e.g. left ventricular outflow obstruction, aortic narrowing) or severe cardiac failure

3. Check concomitant medication use

Please check what other medicines the patient is taking.

If the patient answers **YES** to any of the following or if there is uncertainty around the patient's current medications: **do not supply the product** and refer to the doctor.

- Y N Are you taking nitrates (nicorandil or other nitric oxide donors e.g. glyceryl trinitrate, isosorbide mononitrate or isosorbide dinitrate) for chest pain?
- Y N Are you using drugs called 'poppers' for recreational purposes (e.g. amyl nitrite)?
- Y N Are you taking riociguat or other guanylate cyclase stimulators for lung problems?
- Y N Are you taking ritonavir (for HIV infection)?
- Y N Are you taking any CYP3A4 inhibitors, e.g. saquinavir (to treat HIV infection), cimetidine (a heartburn treatment), itraconazole or ketoconazole (to treat fungal infections), erythromycin or rifampicin (antibiotics) or diltiazem (for high blood pressure)?
- Y N Are you taking any alpha-blockers, such as alfuzosin, doxazosin or tamsulosin, which are medicines to treat urinary problems due to enlarged prostate (benign prostatic hyperplasia) or occasionally to treat high blood pressure?

4. Check concomitant conditions

If the patient answers **YES** to any of the following: **do not supply the product** and refer to the doctor.

- Y N Do you have Peyronie's disease or any other deformation of the penis?
- Y N Have ever had loss of vision because of damage to the optic nerve (such as non-arteritic anterior ischaemic optic neuropathy [NAION]) or have an inherited eye disease (such as retinitis pigmentosa)?
- Y N Do you have galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption?
- Y N Do you have previously diagnosed hepatic (liver) disease (including cirrhosis of the liver) or severe renal (kidney) impairment?
- Y N Do you have any of the following: sickle cell anaemia, multiple myeloma or leukaemia?
- Y N Do you have any bleeding issues (e.g. haemophilia) or have active stomach ulcers?

Additional advice

You should consider possible causes of erectile dysfunction, such as undiagnosed depression, anxiety, excessive alcohol use and taking certain medicines. Examples of classes of medicines that cause ED include diuretics, anti-hypertensives, corticosteroids, anticonvulsants and recreational drugs. Whilst it may be appropriate to supply the product, you should provide lifestyle advice and recommend a follow-up with a doctor.

Dear Doctor,

Please can you review this patient in relation to erectile dysfunction. We had a discussion in my pharmacy, but he was not suitable for supply of Viagra Connect due to his [delete as appropriate] cardiovascular health/interacting medicines/other condition:

.....

Pharmacist signature:..... Date:.....

PHARMACY
STAMP

Please keep this slip and present to the Pharmacist when you next wish to purchase Viagra Connect.

Before resupply your Pharmacist will need to assess you to ensure that there have been no changes in your health or medications since the last time you received the product.

Pharmacist signature:..... Date:.....

PHARMACY
STAMP

Points for counselling and other information

Please follow these points to ensure that your patient is counselled appropriately whether they are supplied the product or not.

Advice for men who have not been supplied Viagra Connect

Men who have not been supplied Viagra Connect because of their cardiovascular health, interacting medicines or another concern must be told to see their GP as soon as they can within 6 months. The tear-off slip (below) can be filled in and given to the man to facilitate his discussion with the doctor.

Advice for men who have been supplied Viagra Connect

Men should be advised:

- They should schedule a health check-up with their doctor as soon as possible within the first 6 months of starting to use Viagra Connect to ensure that their erection problems are not caused by any serious health condition
- Viagra Connect is only intended for men 18 years and older who have erectile dysfunction (ED). Men who do not have ED will not benefit from using this product
- Take one tablet approximately 1 hour before planning to have sexual intercourse. Viagra Connect can start to work within 30 minutes
- Take with or without food, but Viagra Connect may take longer to work after a high-fat meal
- Do not take with grapefruit or grapefruit juice, as it may affect the way the medicine works
- The maximum recommended dosing frequency is one 50 mg tablet per day
- They may need to take Viagra Connect a number of times on different occasions (a maximum of one 50 mg tablet per day), before they can achieve a penile erection satisfactory for sexual activity. If, after several attempts on different dosing occasions, patients are still not able to achieve a penile erection sufficient for satisfactory sexual activity, they should be advised to consult a doctor
- Medicines containing any nitrates (e.g. glyceryl trinitrate, isosorbide mononitrate, isosorbide dinitrate, amyl nitrite also known as 'poppers'), or nitric oxide donors (e.g. sodium nitroprusside or nicorandil), must NOT be used at the same time as Viagra Connect as this combination may lead to a dangerous fall in blood pressure
- Men should tell their doctor that they have started taking Viagra Connect, especially if they are started on any new medicines
- Remind patients about common side effects. These include: headache, flushing, dyspepsia, nasal congestion, dizziness, nausea, visual disturbance, cyanopsia (blue-tinted vision) and blurred vision

Note: If any of these become a concern, advise the patient to talk with a pharmacist or doctor.

Men should be advised to STOP TAKING Viagra Connect and seek medical attention IMMEDIATELY if they experience any of the following SERIOUS side effects. Side effects can be reported by patients or pharmacists via HPRa Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 676 4971; Fax: +353 1 676 2517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects, you can help provide more information on the safety of this medicine.

- Chest pains: If this occurs before, during or after intercourse, they should get into a semi-sitting position and try to relax. Nitrates must NOT be used to treat chest pains
- A persistent and sometimes painful erection lasting longer than 4 hours
- A sudden decrease or loss of vision
- An allergic reaction. Symptoms include sudden wheeziness, difficulty breathing or dizziness, swelling of the eyelids, face, lips or throat
- Serious skin reactions such as Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Syndrome (TEN). Symptoms may include severe peeling and swelling of the skin, blistering of the mouth, genitals and around the eyes, fever
- Seizures or fits

Follow up advice for all men

- ED can be associated with a number of contributing conditions, e.g. hypertension, diabetes mellitus, hypercholesterolaemia or cardiovascular disease. As a result, all men with ED should be advised to consult their doctor within 6 months for a clinical review of potential underlying conditions and risk factors associated with ED
- Provide appropriate advice on lifestyle factors and general healthy living, including:
 - Losing weight
 - Giving up smoking
 - Cutting back on alcohol/recreational drugs
 - Exercising regularly
 - Reducing stress
- You may also want to check if the man is buying products from unregulated sources. It is important to explain these products are not tested for their safety or effectiveness, may not contain the ingredients listed within them and are therefore potentially dangerous, unlike product sourced from a pharmacy and medicines obtained via prescription from the doctor

PP-VCO-IRL-0017 Date of Preparation: October 2020



Your Pharmacist has supplied you with Viagra Connect today for your personal use.

Please make an appointment to see your GP as soon as you can within 6 months of the date of first receiving Viagra Connect (see date on reverse) to check for underlying medical problems that can sometimes be associated with erectile dysfunction.

You have not been provided with Viagra Connect for ED today as it may not be suitable for you to take without consulting your doctor. You should go to see your doctor to discuss other suitable options.

Please make an appointment to see your GP as soon as you can within the next 6 months and present this slip.

Advice for men who are suitable for VIAGRA CONNECT®

Advise patients:

- The product is only for men aged 18 years and over who have ED. Men who do not have ED will not benefit from this product
- To avoid taking nitrates (e.g. glyceryl trinitrate, isosorbide mononitrate, isosorbide dinitrate), nitric oxide donors (e.g. sodium nitroprusside or nicorandil) or amyl nitrite (“poppers”) as these can cause a dangerous drop in blood pressure when used in combination with VIAGRA CONNECT
- During any interaction with their doctor men should tell their doctor that they have started taking VIAGRA CONNECT, especially if they are started on any new medicines
- Men should not take after excessive drinking as VIAGRA CONNECT may be less effective. Excessive alcohol consumption can impact sexual function, as it acts as a central nervous system depressant and can impair motor and cognitive function
- How to take VIAGRA CONNECT:
 - Take 1 tablet approximately 1 hour before planning to have sexual intercourse or to masturbate
 - Swallow the tablet whole, with water
 - Do not take more than 1 tablet per day
 - VIAGRA CONNECT can start to work within 30 minutes, but men are still able to obtain an erection in response to sexual stimulation for up to 4 hours post-dose
 - VIAGRA CONNECT can be taken with or without food but may take longer to work after a high-fat meal
 - Avoid grapefruit juice as this may increase sildenafil levels in the blood
- For most men, VIAGRA CONNECT will work the first or second time they try it. For men who have not been able to have sexual intercourse for some time, it can take additional attempts to obtain maximum benefit. 74% of men will respond to VIAGRA CONNECT (sildenafil 50 mg).¹⁵ If after several attempts (up to 8 separate dosing occasions) on different dosing occasions, patients are still not able to achieve a penile erection sufficient for satisfactory sexual activity, they should be advised to consult their doctor.
- One VIAGRA CONNECT tablet can be taken daily, men should be advised to visit their doctor as soon as possible within 6 months of starting the product for a general health check-up. Once reviewed by their doctor, they can continue using the product, providing there is no change in their circumstances (health or medication).
- If they experience any of the serious side effects detailed in the Side Effect section below, they should stop taking VIAGRA CONNECT and seek IMMEDIATE medical advice.

What if a patient takes more than the recommended dose of VIAGRA CONNECT?

Data show that at doses 16 times the VIAGRA CONNECT dosage, adverse reactions were similar to those seen at lower doses, but the incidence rates and severities were increased. If you are concerned a patient has taken an overdose, refer them immediately to the nearest Hospital Emergency Department or their on-call GP.

Are there any side effects associated with VIAGRA CONNECT?

Sildenafil is generally well tolerated and side effects reported in association with its use are usually transient and mild-to-moderate in intensity.

Patients who experience any of the following after taking VIAGRA CONNECT should be advised to STOP TAKING VIAGRA CONNECT and seek IMMEDIATE medical advice:

- Chest pain, dizziness, feeling of faintness or nausea during or after sex. Nitrates must NOT be used to treat chest pains
- A persistent and sometimes painful erection that lasts more than 4 hours
- A sudden decrease in vision or hearing
- Allergic reaction including sudden wheeziness, difficulty breathing, dizziness or swelling of eyelids, face, lips and throat
- Serious skin reactions which include severe peeling and swelling of the skin, blistering of mouth, genitals and around the eyes, and fever
- Seizure or fits

A full list of side effects experienced with VIAGRA CONNECT can be found in the SmPC which is available on www.medicines.ie.¹⁵

The most commonly reported side effects in the use of VIAGRA CONNECT are:

- Headache
- Facial flushing, hot flush
- Dyspepsia
- Nausea
- Nasal congestion
- Dizziness
- Visual disturbance, blurred vision and blue-tinted vision (cyanopsia)

Advice for men who are not suitable for VIAGRA CONNECT®

- Temporary change in colour vision

Please see Section 2 for men who must not take VIAGRA CONNECT.

If unsuitable for VIAGRA CONNECT, it is important to advise the man of the reason they are not suitable (e.g. it may be a contraindication or that he doesn't have ED). All men with ED should be referred to their GP to explore other suitable options and advised that they should do this as soon as possible within 6 months. Men should be encouraged to seek advice from their doctor for two reasons – because ED may be a symptom of other conditions that need medical attention and because, although VIAGRA CONNECT is not suitable for them, the doctor may be able to prescribe medicines which are suitable.

Advice for all men

ED can be associated with a number of underlying conditions, e.g. hypertension, diabetes mellitus, hypercholesterolaemia or cardiovascular disease. As a result, all men with ED should be advised to consult their doctor within 6 months for a clinical review of potential underlying conditions and risk factors associated with ED.

Lifestyle advice and guidance should be provided to all men, as it can help to reduce the risk factors for ED:

- Weight management
- Healthy eating
- Regular exercise
- Reducing stress
- Quitting smoking
- Moderating alcohol consumption
- Avoiding recreational drugs

As well as helping to improve ED, these changes can also improve general health and may help to reduce the risk of CVD.

Case studies

Case study 3:

How would you respond to the following patient request?

Mr. W is back at your pharmacy having purchased VIAGRA CONNECT for the first time a few days ago. He tells you he's taken a tablet, but it didn't work and he doesn't know why.

How would you handle this conversation? What advice would you give?

See page 35 for case study responses

Case study 4:

How would you respond to the following patient request?

Mr. R is a 58-year-old smoker who, reluctantly, asks if there is anywhere he can speak to you in private. You take him into the consultation area, where you establish he has ED. Mr. R asks to buy 2 packs of VIAGRA CONNECT, but does not want to go into any details about his issue. You ask him how long he has had the problem and he explains that he has been drinking more since he lost his mother last year. He tells you his partner asked him to come to the pharmacy amid worries about his sexual and mental health.

How would you handle this conversation? Would you make a sale of VIAGRA CONNECT in this situation? What other advice could you offer?

Remember: ED may not be the only concern; Mr. R has some signs that could indicate untreated depression

See page 35 for case study responses

Patient supply scenarios

While the decision to supply VIAGRA CONNECT or not will be straightforward in the majority of cases, there will be some situations where the decision is less clear cut. In this section, you will find some less frequent, but more thought-provoking supply scenarios to consider.

Which of the following patients do you feel are suitable for VIAGRA CONNECT.

For some suggested actions see page 31.

- S1** Mr. S is a 57-year-old man and has been a widower for 3 years. He has started a new relationship and is very anxious that this goes well and reveals that he has been having problems with his erectile function for the last 8 months. At his regular health check-up at his doctor 5 months ago, there was nothing untoward found in his assessment. He is in relatively good health and takes ibuprofen for osteoarthritis; he was a smoker (20-a-day) but gave up about 8 years ago.
- S2** Mr. T has had difficulty controlling his hypertension over the last 12 months. After many changes (including following lifestyle advice), he has been stabilised for the past 3 months on a thiazide-related diuretic and an angiotensin-II receptor antagonist combination. Although he is happy with his cardiovascular status, he is having problems with his ability to keep an erection and wonders if VIAGRA CONNECT would be suitable for him.
- S3** Mr. B has been discharged into the care of his GP having had a heart attack 9 months ago that resulted in bypass surgery. After an intense period of cardiac rehabilitation, he is no longer experiencing any chest pain or breathlessness. He explains he is interested in using VIAGRA CONNECT, as he and his wife have missed the physical side of their relationship and his GP has advised him to talk with you about the OTC options available. He is on a standard regimen of drugs including aspirin, a statin and an ACE inhibitor and he also takes a non-steroidal anti-inflammatory drug for osteoarthritis.
- S4** Mr. F is a 36-year-old man and wants a quiet word with you. He is in good health and follows a good diet and exercise regime, but tells you his new girlfriend is hard to satisfy. While he has no issues with having sexual intercourse, he is finding it is taking him longer to recover post coitus and as such wants something to 'pep him up'.
- S5** Mr. M came to see you a few weeks ago for a Medicines Use Review (MUR). During that process, you advised him to talk to his doctor about his prescription for a sublingual glyceryl trinitrate spray. Although he has held a prescription for this drug for a number of years, he has not needed to use this. His MUR shows he is on a standard combination of other drugs to control his symptoms, and as a result of your intervention, his doctor has removed the nitrate from his drug regimen. He is now back and is keen to discuss whether or not VIAGRA CONNECT would be suitable for him as he is experiencing erectile dysfunction. You see from his drug records there are no other contraindicated medicines and he is well with no overt cardiovascular related symptoms.

SECTION 5: Patient supply scenarios

- S6** Mr. A asks for VIAGRA CONNECT as he has recently found it more difficult to perform sexually. On questioning him, you note he is a diabetic and takes a number of tablets for the condition. He also tells you he likes a drink and has been using alcohol more frequently of late. The diabetic nurse has warned him that he may need to start using insulin as his sugar control is very poor.
- S7** Mr. Y has been coming to your pharmacy on a regular basis and has been purchasing VIAGRA CONNECT. In fact, it is 7 months since his first purchase and although you have repeatedly reminded him he needs to follow up with his doctor, he hasn't done so. He has promised he will make an appointment right after he finishes in the pharmacy and tells you it has kept slipping his mind.
- S8** Mr. K came into your pharmacy last week and after taking him through the VIAGRA CONNECT Pharmacy Consultation Guide you supplied an 8-tablet pack of VIAGRA CONNECT. He is back this week and has asked for a further supply of 8 tablets.
- S9** Mr. L is a regular patient who was started on a course of treatment with an SSRI for depression about 6 weeks ago by his GP. Whilst Mr. L is feeling much happier, he has noticed he has had trouble getting and keeping an erection. He is keen to try VIAGRA CONNECT.
- S10** Mr. D is a 70-year-old gentleman who has recently started a new relationship. He tells you he is concerned about the quality of his erection and that he has difficulties maintaining this during sexual intercourse. You ask him about his medication history and he tells you he takes a product for BPH. On further questioning you ascertain the drug is finasteride, which he has been taking for 18 months. He has not had sex since he lost his wife 3 years ago, and he is keen to try VIAGRA CONNECT.
- S11** Mr J is a 65-year-old man who is interested in purchasing VIAGRA CONNECT. You take him through the Pharmacy Consultation Guide and when it comes to questions about his cardiovascular status, all you can ascertain from him is that he has some problems with his heart. As he is a regular customer you check his patient medication record and find he is prescribed the following in the last 12 months: Atenolol 100 mg, Hydrochlorothiazide 25 mg and 2.5 mg amiloride. What advice would you give him?

Patient scenarios: Suggested actions

- S1** VIAGRA CONNECT can be supplied to Mr. S. From his history, the use of the Pharmacy Consultation Guide which has helped ascertain his low CV risk and fitness for sex as well as your professional judgement, he would appear to be in good health. He has given up smoking and he regularly engages with his doctor. The information he has provided helps with the supply decision. For most men, VIAGRA CONNECT will work the first or second time they try it. For men who have not been able to have sexual intercourse for some time, such as Mr. S, it can take additional attempts to obtain maximum benefit (up to 8 separate dosing occasions). If he doesn't respond after 8 doses, he should be advised to consult with his GP. He should also inform his GP he is using the product when he next visits and this consultation should be as soon as possible within 6 months.
- S2** VIAGRA CONNECT can be supplied to Mr. T as his hypertension is now controlled and he is not taking any contraindicated medicines. As an addition to using the Pharmacy Consultation Guide and your professional judgement, it is clear that Mr. T is under the care of a doctor as he has been stabilised with respect to his hypertension. Given what Mr. T has told you it looks like he is fit for sex. Advise Mr. T of the likely side effects of VIAGRA CONNECT and get him to mention he is using the product to his doctor and that he should visit the doctor as soon as possible within 6 months. Advise him to come back if he has any problems or if there are any changes to his health or medication.
- S3** VIAGRA CONNECT can be supplied to Mr. B as it has been more than 6 months since his heart attack and he is not taking any contraindicated medications. Provide Mr. B with the usual follow-up advice. The Pharmacy Consultation Guide has meant that the CV health and fitness for sex has been evaluated and in this case the patient has shown very good signs of improvement in relation to his cardiovascular and general health. He has obviously followed his specialist's advice and has done well with his cardiac rehabilitation. It is important for Mr. B to mention he is using the product to his doctor and as usual he should be counselled on using the product correctly and to go back to the doctor as soon as possible within 6 months.
- S4** VIAGRA CONNECT cannot be supplied to Mr. F as it does not help with shortening the refractory period in sexual intercourse. The drug is only indicated for men who have ED and have described their symptoms in line with the definition of the condition. The no supply slip should be completed and he should be advised to visit his doctor for further advice.
- S5** VIAGRA CONNECT can be supplied to Mr. M. Following the questions on the Pharmacy Consultation Guide, it has been ascertained that he is not taking any contraindicated medications and had no other overt CV issues. As the patient's GP is happy to remove his nitrate prescription and he seems in otherwise good health, he looks like a suitable candidate for the product. The Pharmacy Consultation Guide should be used to provide advice to counsel Mr. M in using the product correctly and advise that he return to see his doctor as soon as possible within 6 months.

SECTION 5: Patient supply scenarios

- S6** VIAGRA CONNECT can be supplied to Mr. A. It is important that he is taken through the Pharmacy Consultation Guide especially with a view on fitness for sex. It appears that he can be supplied the product, but in Mr. A's case there are other issues that warrant investigation, e.g. his increased alcohol use could be a symptom of depression, and he should be advised to speak with his GP and/or his diabetic nurse as soon as possible within 6 months.
- S7** There are no restrictions within the SPC to prevent supply of the product, so the product can be supplied. However, the man should again be reminded that it is very important that he should see his GP and that this may be because there are underlying conditions which may be causing the ED and you may wish to elaborate on these further to help facilitate the man to see his GP. If you get to the point where you feel that the man has not taken your advice, it is longer than 6 months post supply and the man has continued not to see his GP, you are free to use your professional judgement and state you do not wish to supply until you are happy he has been assessed by his GP.
- S8** Provided there has been no change since Mr. K completed the VIAGRA CONNECT Pharmacy Consultation Guide, you can provide a further supply of VIAGRA CONNECT. Remind him the maximum dose is 1 tablet per day. It is also important to check that the drug is working for Mr. K and that he is deriving the benefit he needs. The message to see his GP as soon as possible within 6 months should be repeated.
- S9** Provide Mr. L with positive encouragement that it is good to hear that the anti-depressant is working and that he is feeling happier and reiterate that despite this issue he should not stop taking the anti-depressants. State that ED can be a side effect of anti-depressants, but reinforce that VIAGRA CONNECT can be supplied. Make sure to advise Mr. L to tell his GP he has started taking VIAGRA CONNECT and that he has experienced ED. You should also counsel him on what he should do if VIAGRA CONNECT makes no difference and that in this circumstance he should not discontinue his SSRI medication but return to the GP immediately for further advice.
- S10** VIAGRA CONNECT can be supplied to Mr. D as the Pharmacy Consultation Guide has meant that you have ascertained he is fit for sex, he has good cardiovascular health and is not taking any contraindicated medicines. Provide advice on using the product, particularly as he has not had sex for 3 years, it may take several attempts before it works. For most men, VIAGRA CONNECT will work the first or second time they try it. For men who have not been able to have sexual intercourse for some time, it can take additional attempts to obtain maximum benefit (up to 8 separate dosing occasions). If it doesn't work after 8 different dosing occasions (a maximum of one 50 mg tablet per day) he should see his GP. He should also be advised to see the GP as soon as possible within 6 months of taking VIAGRA CONNECT for a health check.

S11

Mr J's medications could be causing erectile dysfunction as a side effect, and they are not contraindicated with VIAGRA CONNECT. His medications may also give an indication of the conditions he may have. He could be taking atenolol for hypertension, angina or cardiac arrhythmias, and the hydrochlorothiazide/amiloride combination for hypertension or congestive heart failure. Given he has had the same prescriptions for the past 12 months, it is likely that his blood pressure is stable, but you won't be able to know that all other conditions are controlled. It is important to check the patient's CV health status, including fitness for sex, as per the Pharmacy Consultation Guide, as this may not be clear from his medications alone.

If following further evaluation using the Pharmacy Consultation Guide, you decide that Mr J is suitable for VIAGRA CONNECT, it will be important to advise Mr J on how to use the product. If the CV health status and suitability cannot be deduced from your discussions with the man, then it is probably best to refuse supply and refer him to his doctor.

Case Study Responses

Case study 1

Given that ED can be an early sign of the development of underlying CVD and coupled with the fact Mr. P is getting out of breath quickly, it is important that he is advised to visit his doctor for a check-up. There is an opportunity to discuss Mr. P's lifestyle and provide him with some information around this, but it is important he is not supplied the product and advised to take further advice from his GP. This guide and the Pharmacy Consultation Guide provide more information around these issues.

Case study 2

Mr. G falls in the category of patients using nitrates and in this case you must not supply VIAGRA CONNECT to him. Taking a nitrate and VIAGRA CONNECT together may lead to a significant drop in blood pressure. You should advise Mr. G of this fact and advise him to speak to his GP about this. In addition, it is important to provide counselling around his other medical issues and again ask him to speak to his doctor about his current health status. You can also offer him advice around his lifestyle choices.

Case study 3

For some patients VIAGRA CONNECT may not work on taking the first few doses. This can be for a variety of reasons; the man may have been out of practice for a while or may need a higher dose or an alternative treatment. With Mr. W, it is important to establish, how he used the product, e.g. did he take it on an empty stomach or after a meal? Did he take it one hour before he planned to have sexual intercourse?

Once correct use is established and advised on, it will be important to understand what his situation is. For example, is this a new relationship or an existing one? Was he nervous because he hadn't had sex for some time? Experts advise men may need to take up to 8 doses of a PDE5i on different occasions before they are considered a non-responder. In this case it may be worth advising Mr. W to try the product on his own to get used to the effects before proceeding to sexual intercourse. If after a few attempts the product is still not providing a satisfactory response, Mr. W should be advised to come back to speak to you or to visit his GP for further advice.

Case study 4

Mr. R should be taken through the Pharmacy Consultation Guide to establish his symptoms and to look at the underlying issues. You know he is a smoker and there is a possibility that he is not in the best of health. The Pharmacy Consultation Guide is designed to help you explore these issues and ascertain his suitability for the product. It is important to remember ED may not be the only concern; Mr. R has some signs that could indicate untreated depression. As such Mr. R should be advised to see his doctor to get a thorough assessment of his physical and mental health. The decision to supply the product will depend on the answers he gives you and this guide will help you with determining this for the wide range of men you may need to talk to.

Essential information

Name of product: VIAGRA CONNECT 50 mg film-coated tablets.

Active ingredient: sildenafil.

Further information is available in the SmPC.

Supply classification: Pharmacy only.

Indications: For erectile dysfunction in adult men.

Dosage and Method of use: Adults: The recommended dose is one 50 mg tablet taken with water approximately one hour before sexual activity. The maximum recommended dosing frequency is once per day. If Viagra CONNECT is taken with food, the onset of activity may be delayed compared to the fasted state. Patients should be advised that they may need to take Viagra CONNECT a number of times on different occasions (a maximum of one 50 mg tablet per day), before they can achieve a penile erection satisfactory for sexual activity. If after several attempts on different dosing occasions patients are still not able to achieve a penile erection sufficient for satisfactory sexual activity, they should be advised to consult a doctor. Elderly: Dosage adjustments are not required in elderly patients (≥ 65 years old). Renal Impairment: No dosage adjustments are required for patients with mild to moderate renal impairment. However, since sildenafil clearance is reduced in individuals with severe renal impairment (creatinine clearance <30 ml/min), individuals previously diagnosed with severe renal impairment must be advised to consult their doctor before taking Viagra CONNECT, since a 25 mg tablet may be more suitable for them (see Warnings and Precautions). Hepatic Impairment: Sildenafil clearance is reduced in individuals with hepatic impairment (e.g. cirrhosis). Individuals previously diagnosed with mild to moderate hepatic impairment must be advised to consult their doctor before taking Viagra CONNECT, since a 25 mg tablet may be more suitable for them (see Warnings and Precautions). The safety of sildenafil has not been studied in patients with severe hepatic impairment, and its use is therefore contraindicated (see Contra-indications). Use in patientstaking other medicinal products: Pharmacokinetic analysis of clinical trial data indicated a reduction in sildenafil clearance when co-administered with CYP3A4 inhibitors (such as ritonavir, ketoconazole, itraconazole, erythromycin, cimetidine). With the exception of ritonavir, for which co-administration with sildenafil is contraindicated (see Contra-indications), individuals receiving concomitant treatment with CYP3A4 inhibitors must be advised to consult their doctor before taking Viagra CONNECT, since a 25 mg tablet may be more suitable for them (see Warnings and Precautions). In order to minimise the potential of developing postural hypotension in patients receiving alpha blocker treatment (e.g. alfuzosin, doxazosin or tamsulosin), patients should be stabilised on alpha blocker therapy prior to initiating sildenafil treatment. Thus, patients taking alpha blockers must be advised to consult their doctor before taking Viagra CONNECT since a 25 mg tablet may be more suitable for them (see Warnings and Precautions).

Side Effects: The most commonly reported adverse reactions in clinical studies among sildenafil treated patients were headache, flushing, dyspepsia, nasal congestion, dizziness, nausea, hot flush, visual disturbance, cyanopsia and vision blurred. Adverse reactions from post-marketing surveillance has been gathered covering an estimated period >10 years. Because not all adverse reactions are reported to the Marketing Authorisation Holder and included in the safety database, the frequencies of these reactions cannot be reliably determined. Very Common ($\geq 1/10$): Headache. Common ($\geq 1/100$ and $<1/10$): Dizziness, Visual colour distortions (Chloropsia, Chromatopsia, Cyanopsia, Erythroptopsia and Xanthopsia), Visual disturbance, Vision blurred, Flushing, Hot flush, Nasal congestion, Nausea, Dyspepsia. Uncommon ($\geq 1/1,000$ and $<1/100$): Rhinitis, Hypersensitivity; Somnolence; Hypoaesthesia, Lacrimation disorders (Dry eye, Lacrimal disorder and Lacrimation increased), Eye pain, Photophobia, Photopsia, Ocular hyperaemia, Visual brightness, Conjunctivitis, Vertigo, Tinnitus, Tachycardia, Palpitations, Hypertension, Hypotension, Epistaxis, Sinus congestion, Gastro Oesophagael reflux disease, Vomiting, Abdominal pain upper, Dry mouth, Rash, Myalgia, Pain in extremity, Haematuria, Chest pain, Fatigue, Feeling hot, Heart rate increased. Rare ($\geq 1/10,000$ and $<1/1,000$): Cerebrovascular accident, Transient ischaemic attack, Seizure, Seizure recurrence, Syncope, Non-arteritic anterior ischaemic optic neuropathy (NAION), Retinal vascular occlusion, Retinal haemorrhage, Arteriosclerotic retinopathy, Retinal disorder, Glaucoma, Visual field defect, Diplopia, Visual acuity reduced, Myopia, Asthenopia, Vitreous floaters, Iris disorder, Mydriasis, Halo vision, Eye oedema, Eye swelling, Eye disorder, Conjunctival hyperaemia, Eye irritation, Abnormal sensation in eye, Eyelid oedema, Scleral discoloration, Deafness, Sudden cardiac death, Myocardial infarction, Ventricular arrhythmia, Atrial fibrillation, Unstable angina, Throat tightness, Nasal oedema, Nasal dryness, Hypoaesthesia oral, Stevens-Johnson Syndrome (SJS), Toxic Epidermal Necrolysis (TEN), Penile haemorrhage, Priapism, Haemospermia, Erection increased, Irritability.

Warnings and Precautions: Erectile dysfunction can be associated with a number of contributing conditions, e.g. hypertension, diabetes mellitus, hypercholesterolaemia or cardiovascular disease. As a result, all men with erectile dysfunction should be advised to consult their doctor within 6 months for a clinical review of potential underlying conditions and risk factors associated with erectile dysfunction (ED). If symptoms of ED have not improved after taking Viagra CONNECT on several consecutive occasions, or if their erectile dysfunction worsens, the patient should be advised to consult their doctor. Cardiovascular risk factors: Since there is a degree of cardiac risk associated with sexual activity, the cardiovascular status of men should be considered prior to initiation of therapy. Agents for the treatment of erectile dysfunction, including sildenafil, are not recommended to be used by those men who with light or moderate physical activity, such as walking briskly for 20 minutes or climbing 2 flights of stairs, feel very breathless or experience chest pain. The following patients are considered at low cardiovascular risk from sexual activity: patients who have been successfully revascularised (e.g. via coronary artery bypass grafting, stenting, or angioplasty), patients with asymptomatic controlled hypertension, and those with mild valvular disease. These patients may be suitable for treatment but should consult a doctor before resuming sexual activity. Patients previously diagnosed with the following must be advised to consult with their doctor before resuming sexual activity: uncontrolled hypertension, moderate to severe valvular disease, left ventricular dysfunction, hypertrophic obstructive and other cardiomyopathies, or significant arrhythmias. Sildenafil has vasodilator properties, resulting in mild and transient decreases in blood pressure. Patients with increased susceptibility to vasodilators include those with left ventricular outflow obstruction (e.g. aortic stenosis), or those with the rare syndrome of multiple system atrophy manifesting as severely impaired autonomic control of blood pressure. Men with these conditions must not use the product without consulting a doctor. Sildenafil potentiates the hypotensive effect of nitrates (see Contra- indications). Serious cardiovascular events, including myocardial infarction, unstable angina, sudden cardiac death, ventricular arrhythmia, cerebrovascular haemorrhage, transient ischaemic attack, hypertension and hypotension have been reported post-marketing in temporal association with the use of sildenafil. Most, but not all, of these patients had pre-existing cardiovascular risk factors. Many events were reported to occur during or shortly after sexual intercourse and a few were reported to occur shortly after the use of sildenafil without sexual activity. It is not possible to determine whether these events are related directly to these factors or to other factors. Priapism: Patients who have conditions which may predispose them to priapism (such as sickle cell anaemia, multiple myeloma or leukaemia), should consult a doctor before using agents for the treatment of erectile dysfunction, including sildenafil. Prolonged erections and priapism have been occasionally reported with sildenafil in post-marketing experience. In the event of an erection that persists longer than 4 hours, the patient should seek immediate medical assistance. If priapism is not treated immediately, penile tissue damage and permanent loss of potency could result. Concomitant use with other treatments for erectile dysfunction: The safety and efficacy of combinations of sildenafil with other treatments for erectile dysfunction have not been studied. Therefore the use of such combinations is not recommended. Effects on vision: Cases of visual defects have been reported spontaneously in connection with the intake of sildenafil and other PDE5 inhibitors (see Side Effects). Cases of non-arteritic anterior ischaemic optic neuropathy, a rare condition, have been reported spontaneously and in an observational study in connection with the intake of sildenafil and other PDE5 inhibitors (see Side Effects). Patients should be advised that in the event of any sudden visual defect, they should stop taking Viagra CONNECT and consult a physician immediately. Concomitant use with CYP3A4 inhibitors: Pharmacokinetic analysis of clinical trial data indicated a reduction in sildenafil clearance when co-administered with CYP3A4 inhibitors (such as ketoconazole, itraconazole, erythromycin, cimetidine). Although, no increased incidence of adverse events was observed in these patients, they should be advised to consult a doctor before taking Viagra CONNECT as a 25 mg tablet may be more suitable for them. Concomitant use with alpha-blockers: Caution is advised when sildenafil is administered to patients taking an alpha-blocker, as the co-administration may lead to symptomatic hypotension in a few susceptible individuals. This is most likely to occur within 4 hours post sildenafil dosing. In order to minimise the potential for developing postural hypotension, patients should be hemodynamically stable on alpha-blocker therapy prior to initiating sildenafil treatment. Thus, patients taking alpha blockers should be advised to consult their doctor before taking Viagra CONNECT as a 25 mg tablet may be more suitable for them. Treatment should be stopped if symptoms of postural hypotension occur, and patients should seek advice from their doctor on what to do. Effect on bleeding: Studies with human platelets indicate that sildenafil potentiates the antiaggregatory effect of sodium nitroprusside in vitro. There is no safety information on the administration of sildenafil to patients with bleeding disorders or active peptic ulceration. Therefore the use of sildenafil is not recommended in those patients with history of bleeding disorders or active peptic ulceration, and should only be administered after consultation with a doctor. Hepatic impairment: Patients with hepatic impairment must be advised to consult their doctor before taking Viagra CONNECT, since a 25 mg tablet may be more suitable for them. Renal impairment: Patients with severe renal impairment

(creatinine clearance <30 mL/min), must be advised to consult their doctor before taking Viagra CONNECT, since a 25 mg tablet may be more suitable for them. Lactose: The film coating of the tablet contains lactose. Viagra CONNECT should not be administered to men with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption. Sodium: This medicinal product contains less than 1 mmol sodium (23 mg) per tablet. Patients on low sodium diets can be informed that this medicinal product is essentially 'sodium-free'. Use with alcohol: Drinking excessive alcohol can temporarily reduce a man's ability to get an erection. Men should be advised not to drink large amounts of alcohol before sexual activity.

Contra-indications: Hypersensitivity to the active substance or to any of the excipients. Consistent with its known effects on the nitric oxide/cyclic guanosine monophosphate (cGMP) pathway, sildenafil was shown to potentiate the hypotensive effects of nitrates, and its co-administration with nitric oxide donors (such as amyl nitrite) or nitrates in any form is therefore contraindicated. Co-administration of Viagra CONNECT with ritonavir (a highly potent P450 enzyme inhibitor) is contraindicated. The co-administration of phosphodiesterase type 5 (PDE5) inhibitors, including sildenafil, with guanylate cyclase stimulators, such as riociguat, is contraindicated as it may potentially lead to symptomatic hypotension. Agents for the treatment of erectile dysfunction, including sildenafil, should not be used by those men for whom sexual activity may be inadvisable, and these patients should be referred to their doctor. This includes patients with severe cardiovascular disorders such as a recent (6 months) acute myocardial infarction (AMI) or stroke, unstable angina or severe cardiac failure. Sildenafil should not be used in patients with severe hepatic impairment, hypotension (blood pressure < 90/50 mmHg) and known hereditary degenerative retinal disorders such as *retinitis pigmentosa* (a minority of these patients have genetic disorders of retinal phosphodiesterases). This is because the safety of sildenafil has not been studied in these sub-groups of patients, and its use is therefore contraindicated. Sildenafil is contraindicated in patients who have loss of vision in one eye because of non-arteritic anterior ischaemic optic neuropathy (NAION), regardless of whether this episode was in connection or not with previous PDE5 inhibitor exposure. Viagra CONNECT should not be used in patients with anatomical deformation of the penis (such as angulation, cavernosal fibrosis or Peyronie's disease). Viagra CONNECT is not indicated for use by women. The product is not intended for men without erectile dysfunction. This product is not intended for men under 18 years of age.

Legal Category: Product not subject to medical prescription.

Marketing Authorisation Number: PA0822/234/001.

Marketing Authorisation Holder: Pfizer Healthcare Ireland, 9 Riverwalk, Citywest Business Campus, Dublin 24, Ireland.

For further information on this medicine please contact: Pfizer Medical Information on 1800 633 363 or at EUMEDINFO@pfizer.com. For queries regarding product availability please contact: Pfizer Healthcare Ireland, Pfizer Building 9, Riverwalk, National Digital Park, Citywest Business Campus, Dublin 24 + 353 1 467 6500.

Date: 10/2020.

Ref: VC 1_0.

References

1. Lue T, *et al.* Summary of the recommendations on sexual dysfunctions in men. *J Sex Med* 2004;1:6-23.
2. Kubin M, Wagner G, Fugl-Meyer AR. Epidemiology of erectile dysfunction. *Int J Impot Res*. 2003;15:63-71 .
3. Miner M. Erectile dysfunction: a harbinger or consequence: does its detection lead to a window of curability? *J Androl* 2011;32:125-134.
4. The British Association of Urological Surgeons. Erectile dysfunction. Frequently asked questions. Available at: https://www.baus.org.uk/patients/conditions/3/erectile_dysfunction_impotence. Accessed Nov 2017.
5. Rosen R. Psychogenic erectile dysfunction: classification and management. *Urol Clin North Am* 2001;28:269-278.
6. Conaglen H, *et al.* Drug-induced sexual dysfunction in men and women. *Aust Prescr* 2013;36:42-52.
7. Nunes KP, *et al.* New insights into hypertension-associated erectile dysfunction. *Curr Opin Nephrol Hypertens* 2012;21:163-170.
8. Fusco F, *et al.* The impact of non-urologic drugs on sexual function in men. *Arch Ital Urol Androl* 2014;86:1.
9. Lee JC. Club drugs' and erectile function: far from sexual ecstasy. *J Sex Reprod Med* 2002;2:28-30.
10. Lasker GF, *et al.* A review of the pathophysiology and novel treatments for erectile dysfunction. *Adv Pharmacol Sci* 2010;2010:1-10.
11. WebMD. Drugs linked to erectile dysfunction. Available at: <http://www.webmd.com/erectile-dysfunction/guide/drugs-linked-erectile-dysfunction>. Accessed Jun 2017.
12. Muneer A, *et al.* Erectile dysfunction. *BMJ* 2014;348:g129.
13. Cappelleri JC, *et al.* Clinically meaningful improvement on the self-esteem and relationship questionnaire in men with erectile dysfunction. *Qual Life Res* 2007;16:1203-1210.
14. Ballard SA, *et al.* Effects of sildenafil on the relaxation of human corpus cavernosum tissue *in vitro* and on the activities of cyclic nucleotide phosphodiesterase isozymes. *J Urol* 1998;159:2164-2171.
15. VIAGRA CONNECT 50 mg film-coated tablets (sildenafil): Summary of Product Characteristics. Pfizer Healthcare Ireland.
16. The European Association of Urology Guidelines. Available at: https://uroweb.org/wp-content/uploads/Guidelines_WebVersion_Complete-1.pdf. Accessed May 2020.
17. Nehra A, *et al.* The Princeton III consensus recommendations for the management of erectile dysfunction and cardiovascular disease. *Mayo Clin Proc* 2012;87:766-778.

